
**INTERNATIONAL JOURNAL OF CURRENT RESEARCH IN
CHEMISTRY AND PHARMACEUTICAL SCIENCES**

(p-ISSN: 2348-5213; e-ISSN: 2348-5221)

www.ijcrps.com

(A Peer Reviewed, Referred, Indexed and Open Access Journal)

DOI: 10.22192/ijcrps

Coden: IJCROO(USA)

Volume 11, Issue 9- 2024

Review Article



DOI: <http://dx.doi.org/10.22192/ijcrps.2024.11.09.004>

Antioxidant Supplementation and Prevention of Early Pregnancy Loss: A Narrative Review

***Emmanuel Ifeanyi Obeagu¹ and Getrude Uzoma Obeagu²**

¹Department of Medical Laboratory Science, Kampala International University, Uganda.

²School of Nursing Science, Kampala International University, Uganda.

*Corresponding author: Emmanuel Ifeanyi Obeagu, Department of Medical Laboratory Science, Kampala International University, Uganda, emmanuelobeagu@yahoo.com,
ORCID: 0000-0002-4538-0161

Copyright © 2024. Emmanuel Ifeanyi Obeagu and Getrude Uzoma Obeagu. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Early pregnancy loss (EPL), commonly known as miscarriage, affects a significant percentage of pregnancies and poses a considerable emotional burden on individuals and couples. Recent studies have indicated that oxidative stress plays a pivotal role in the pathophysiology of EPL by disrupting cellular function and impairing reproductive health. An imbalance between reactive oxygen species (ROS) and antioxidants can compromise oocyte quality, embryo development, and uterine receptivity, contributing to an increased risk of miscarriage. Antioxidant supplementation has emerged as a promising strategy to mitigate oxidative stress and improve reproductive outcomes. Various antioxidants, including vitamins C and E, Coenzyme Q10, N-acetylcysteine, and omega-3 fatty acids, have demonstrated beneficial effects in enhancing fertility and supporting early pregnancy. This narrative review aims to explore the mechanisms of oxidative stress related to EPL, examine the role of antioxidant defense systems, and present clinical evidence supporting antioxidant supplementation as a preventive measure against EPL.

Keywords: Antioxidant supplementation, early pregnancy loss, oxidative stress, miscarriage, reproductive health

Introduction

Early pregnancy loss (EPL), defined as the spontaneous loss of a pregnancy before the 20th week, is a common occurrence that affects an estimated 10-20% of clinically recognized pregnancies. This percentage may be higher, as many biochemical pregnancies, which are often undetected, also result in loss before a missed period. EPL can cause significant emotional and psychological distress, impacting the well-being of individuals and couples attempting to conceive. One of the critical factors implicated in EPL is oxidative stress, a condition characterized by an imbalance between the production of reactive oxygen species (ROS) and the body's antioxidant defenses. In normal physiological conditions, ROS play a vital role in cellular signaling and homeostasis. However, excessive ROS levels can lead to cellular damage, inflammation, and apoptosis, adversely affecting reproductive health. In the context of early pregnancy, oxidative stress can impair oocyte quality, disrupt embryo development, and hinder uterine receptivity, all of which are essential for a successful pregnancy.¹⁻⁵ The reproductive system is particularly vulnerable to oxidative stress due to its high metabolic activity and the processes involved in gametogenesis and embryo implantation. Factors such as advanced maternal age, environmental toxins, poor nutrition, and underlying health conditions can exacerbate oxidative stress levels, further increasing the risk of EPL. Given the significant role that oxidative stress plays in reproductive failure, there is a growing interest in exploring potential interventions to mitigate its effects, particularly through antioxidant supplementation. Antioxidant supplementation involves providing the body with additional antioxidants to counteract oxidative stress and improve overall reproductive health. Various antioxidants, including vitamins C and E, Coenzyme Q10, N-acetylcysteine (NAC), and omega-3 fatty acids, have been studied for their potential to enhance fertility and support early pregnancy. These compounds work by neutralizing ROS, reducing inflammation, and promoting cellular health, thereby creating a more favorable environment for embryo development

and implantation.⁶⁻¹⁰ Recent clinical studies have begun to investigate the efficacy of antioxidant supplementation in preventing EPL. Evidence suggests that antioxidants may improve oocyte quality, increase implantation rates, and reduce the incidence of miscarriage in certain populations, particularly women undergoing assisted reproductive technologies (ART) or those with known risk factors for EPL. However, despite promising findings, the current body of literature remains limited, and further research is necessary to establish clear guidelines regarding the use of antioxidants in clinical practice.¹¹⁻¹²

Mechanisms of Oxidative Stress in Early Pregnancy Loss

Oxidative stress arises when there is an imbalance between reactive oxygen species (ROS) production and the body's antioxidant defenses, leading to cellular damage and dysfunction. In the context of early pregnancy loss (EPL), this phenomenon can significantly disrupt key physiological processes essential for successful conception and embryo development. Several mechanisms illustrate how oxidative stress can impact early pregnancy and contribute to EPL.¹³⁻¹⁴

- 1. Oocyte Quality and Maturation:** The quality of the oocyte is crucial for fertilization and early embryonic development. During oocyte maturation, an increase in metabolic activity results in elevated ROS levels. If antioxidant defenses are insufficient to counteract this surge, oxidative damage can occur, negatively affecting the oocyte's genetic material and subsequent embryo quality. Studies have shown that poor oocyte quality is associated with a higher risk of miscarriage, highlighting the importance of maintaining redox balance during this critical phase.¹⁴⁻¹⁵

- 2. Fertilization and Embryo Development:** Once fertilization occurs, the embryo undergoes rapid cell divisions and differentiation. High levels of ROS can disrupt these processes, leading to impaired embryo development and viability. Oxidative stress can affect key cellular pathways,

such as mitochondrial function, which is essential for energy production during early embryogenesis. Mitochondrial dysfunction can result in increased apoptosis and cell death, further contributing to the risk of EPL.¹⁶⁻¹⁷

3. Uterine Environment: The uterine environment plays a vital role in embryo implantation and maintenance of early pregnancy. Oxidative stress can alter the uterine microenvironment by promoting inflammation and affecting endometrial receptivity. Elevated ROS levels may lead to increased expression of inflammatory cytokines and chemokines, which can disrupt the normal implantation process and compromise uterine health. A suboptimal uterine environment can hinder successful embryo implantation, increasing the likelihood of EPL.¹⁸⁻¹⁹

4. Placental Development: The placenta is crucial for nutrient and oxygen exchange between the mother and fetus. Proper placentation requires a finely tuned balance of ROS and antioxidants. Excessive oxidative stress can disrupt placental trophoblast function, leading to inadequate remodeling of maternal blood vessels and reduced blood flow. This can result in poor placental development, negatively impacting fetal growth and increasing the risk of EPL.²⁰⁻²¹

5. Genetic and Epigenetic Alterations: Oxidative stress can induce genetic mutations and epigenetic modifications in oocytes and embryos. These changes can lead to impaired cellular signaling and abnormal gene expression, contributing to developmental defects and reproductive failure. For instance, oxidative DNA damage may result in mutations that disrupt normal embryonic development, ultimately leading to miscarriage.²²

6. Hormonal Regulation: Hormones play a critical role in maintaining early pregnancy, and oxidative stress can disrupt hormonal balance. For example, high ROS levels can impair the function of luteal cells, which are responsible for

producing progesterone—a hormone essential for sustaining pregnancy. Insufficient progesterone levels can lead to luteal phase defects, compromising the uterine lining's ability to support embryo implantation and increasing the risk of EPL.²³⁻²⁴

7. Immunological Factors: The immune system plays a vital role in pregnancy by promoting tolerance towards the developing embryo. Oxidative stress can alter immune cell function and cytokine profiles, leading to an inflammatory environment that is detrimental to pregnancy. An imbalance in immune responses may result in maternal immune activation against the embryo, further contributing to the risk of EPL.²⁵⁻²⁶

Antioxidant Defense Systems

The body employs a complex and efficient antioxidant defense system to combat oxidative stress and maintain redox homeostasis. This system comprises both enzymatic and non-enzymatic antioxidants that work synergistically to neutralize reactive oxygen species (ROS) and mitigate their harmful effects on cellular structures. Understanding these defense mechanisms is crucial for recognizing how they can be supported or enhanced, especially in the context of early pregnancy loss (EPL).²⁷⁻²⁸

1. Enzymatic Antioxidants: These are proteins that catalyze biochemical reactions to convert ROS into less harmful substances. Key enzymatic antioxidants include:

- **Superoxide Dismutase (SOD):** SOD is one of the first lines of defense against oxidative stress, catalyzing the conversion of superoxide radicals into hydrogen peroxide. There are different forms of SOD, including cytosolic (Cu/Zn-SOD) and mitochondrial (Mn-SOD), each playing a vital role in specific cellular compartments.²⁹

- **Catalase:** This enzyme further detoxifies hydrogen peroxide generated by SOD, converting it into water and oxygen. Catalase is primarily located in peroxisomes and is crucial for

preventing the accumulation of hydrogen peroxide, which can be harmful in high concentrations.³⁰

○ **Glutathione Peroxidase (GPx):** GPx utilizes glutathione (GSH), a tripeptide antioxidant, to reduce hydrogen peroxide and lipid peroxides, thereby protecting cellular membranes from oxidative damage. GPx plays a critical role in maintaining cellular redox balance and ensuring proper cellular function.³¹

2. **Non-Enzymatic Antioxidants:** These are small molecules that scavenge ROS directly or enhance the effectiveness of enzymatic antioxidants. Important non-enzymatic antioxidants include:

○ **Glutathione:** GSH is a key intracellular antioxidant that plays a vital role in detoxifying ROS and maintaining cellular homeostasis. It can directly scavenge free radicals and also serves as a substrate for GPx in the reduction of hydrogen peroxide. Glutathione levels can be affected by dietary intake and overall health, emphasizing its importance in supporting reproductive health.³²

○ **Vitamins C and E:** Vitamin C (ascorbic acid) is a potent water-soluble antioxidant that can regenerate vitamin E in its active form. It is crucial for neutralizing ROS in extracellular fluids and protecting tissues from oxidative damage. Vitamin E (tocopherol) is a fat-soluble antioxidant that protects cell membranes from lipid peroxidation by trapping free radicals. Together, these vitamins contribute to a robust antioxidant defense.³³⁻³⁴

○ **Coenzyme Q10 (CoQ10):** This lipid-soluble antioxidant is essential for mitochondrial function and energy production. CoQ10 participates in electron transport and can directly scavenge free radicals, helping to mitigate oxidative stress within cells, particularly in energy-demanding tissues such as those involved in reproduction.³⁵

3. **Dietary Antioxidants:** Certain dietary components can enhance the body's antioxidant defenses. Phytochemicals found in fruits and vegetables, such as flavonoids, polyphenols, and carotenoids, possess significant antioxidant properties. These compounds can scavenge free radicals, reduce oxidative damage, and improve overall antioxidant capacity. A diet rich in antioxidants may promote reproductive health by supporting the body's defense systems.³⁶

4. **Redox Signaling:** While antioxidants primarily serve to neutralize ROS, it is essential to note that low levels of ROS play a role in redox signaling, which regulates various cellular processes. Antioxidants help maintain the delicate balance between oxidative stress and redox signaling, ensuring that ROS levels are adequate for normal physiological functions without causing damage. This balance is particularly critical during early pregnancy, where signaling pathways are involved in implantation and placentation.³⁷⁻³⁸

5. **Hormonal Influence:** Hormones such as estrogen and progesterone have been shown to modulate the expression of antioxidant enzymes and influence the body's overall antioxidant status. For example, estrogen can enhance the activity of SOD and GPx, contributing to a more robust antioxidant defense system. Hormonal fluctuations during the menstrual cycle and early pregnancy can therefore impact oxidative stress levels and the effectiveness of antioxidant defenses.³⁹

6. **Mitochondrial Function:** Mitochondria are both a source of ROS and a critical site for antioxidant activity. They contain their own antioxidant systems, including SOD and GPx, which are essential for protecting mitochondrial function and maintaining cellular energy levels. Healthy mitochondrial function is crucial for oocyte and embryo quality, emphasizing the importance of antioxidants in reproductive health.⁴⁰

7. **Clinical Implications:** Understanding the body's antioxidant defense systems highlights the potential for antioxidant supplementation as a therapeutic approach to reduce oxidative stress and improve reproductive outcomes. For women at risk of early pregnancy loss, enhancing antioxidant capacity through dietary means or targeted supplementation could provide significant benefits, supporting overall reproductive health and mitigating the risk of EPL.⁴¹

Antioxidant Supplementation in Preventing Early Pregnancy Loss

Antioxidant supplementation has emerged as a potential strategy for mitigating oxidative stress and improving reproductive outcomes, particularly in the context of early pregnancy loss (EPL). This approach aims to bolster the body's natural defenses against oxidative damage, which can impair various physiological processes crucial for successful conception and embryonic development. The following sections discuss the rationale for antioxidant supplementation, the types of antioxidants commonly used, and the evidence supporting their efficacy in preventing EPL.⁴²

1. **Rationale for Antioxidant Supplementation:** The pathophysiology of early pregnancy loss is multifaceted, with oxidative stress playing a significant role in disrupting cellular functions essential for maintaining pregnancy. Elevated levels of reactive oxygen species can lead to DNA damage, mitochondrial dysfunction, and inflammatory responses that compromise embryo viability. By supplementing with antioxidants, it is possible to enhance the body's capacity to neutralize excess ROS, thereby reducing the likelihood of oxidative damage to oocytes, embryos, and uterine tissue. This supportive approach is particularly important for women with underlying conditions associated with increased oxidative stress, such as polycystic ovary syndrome (PCOS), obesity, or advanced maternal age.⁴³

2. **Types of Antioxidants Used in Supplementation:** Various antioxidants have been studied for their potential benefits in reproductive health, particularly in preventing EPL. Commonly used antioxidants include:

- **Vitamin C:** As a powerful water-soluble antioxidant, vitamin C protects cellular components from oxidative damage. It plays a critical role in collagen synthesis and hormone regulation, both of which are essential for maintaining a healthy uterine environment. Some studies have suggested that vitamin C supplementation may improve oocyte quality and enhance pregnancy outcomes.⁴⁴

- **Vitamin E:** This fat-soluble antioxidant is known for its ability to prevent lipid peroxidation in cell membranes. Vitamin E supplementation has been associated with improved reproductive outcomes, as it may enhance the quality of oocytes and embryos while also supporting healthy placentation.⁴⁵

- **Coenzyme Q10 (CoQ10):** CoQ10 is essential for mitochondrial function and energy production, making it a valuable supplement for women facing infertility or recurrent pregnancy loss. Studies indicate that CoQ10 supplementation can improve oocyte and embryo quality by enhancing mitochondrial efficiency and reducing oxidative stress.⁴⁶

- **N-acetylcysteine (NAC):** NAC is a precursor to glutathione, one of the body's most important intracellular antioxidants. It has been studied for its potential to improve insulin sensitivity in women with PCOS and may also enhance ovarian function and embryo viability through its antioxidant properties.

3. **Evidence Supporting Antioxidant Supplementation:** A growing body of literature supports the use of antioxidant supplementation in improving reproductive outcomes and preventing early pregnancy loss. For example, several clinical studies have reported positive correlations

between antioxidant intake and improved oocyte quality, higher fertilization rates, and better embryo development.

- **Vitamin C and E:** Research has shown that women who receive vitamin C and E supplementation prior to and during pregnancy exhibit reduced oxidative stress markers and improved pregnancy outcomes. A randomized controlled trial indicated that supplementation with these vitamins decreased the rate of EPL in high-risk populations, emphasizing their protective role in early gestation.⁴⁷

- **CoQ10:** Emerging evidence suggests that CoQ10 supplementation may enhance oocyte quality and improve embryo viability in women undergoing assisted reproductive technologies (ART). Studies have found that women who supplemented with CoQ10 had higher pregnancy rates and lower rates of miscarriage compared to those who did not.⁴⁸

- **NAC:** Clinical trials have demonstrated that NAC supplementation can improve reproductive outcomes in women with PCOS, a condition often associated with increased oxidative stress. The use of NAC has been linked to improved ovarian response and embryo quality, potentially reducing the risk of EPL.⁴⁹

4. **Dosing and Safety Considerations:** The effective dosing of antioxidant supplements is crucial to achieving the desired protective effects without adverse outcomes. While many studies have used varying dosages, clinical guidelines for antioxidant supplementation in reproductive health are still evolving. It is important for healthcare providers to tailor antioxidant supplementation regimens to individual patients, considering factors such as age, underlying health conditions, and specific reproductive goals.⁵⁰

- **Safety:** Generally, antioxidant supplements are considered safe when taken within recommended guidelines. However,

excessive intake of certain antioxidants, particularly in isolated forms, may have pro-oxidative effects or interfere with physiological processes. It is vital to encourage women to consult with healthcare professionals before initiating any supplementation, especially during pregnancy.

5. **Combination Therapy:** The use of combination antioxidant therapies may provide synergistic benefits in preventing early pregnancy loss. For example, a regimen that includes both vitamins C and E, along with CoQ10, could target multiple pathways of oxidative stress and enhance overall antioxidant capacity. Clinical studies investigating the efficacy of multi-antioxidant formulations are needed to determine the optimal combinations and dosages for improving reproductive health.⁵¹

Clinical Evidence Supporting Antioxidant Supplementation

The use of antioxidant supplementation in preventing early pregnancy loss (EPL) is supported by a growing body of clinical evidence that underscores its potential to mitigate oxidative stress and improve reproductive outcomes. This section reviews the key studies and findings that highlight the efficacy of various antioxidants in enhancing fertility and reducing the risk of EPL.

1. **Vitamin C and E Supplementation:** Several studies have examined the impact of vitamins C and E on reproductive health. A notable randomized controlled trial involving women with a history of recurrent pregnancy loss found that supplementation with vitamins C and E significantly reduced oxidative stress markers and improved pregnancy outcomes. Women who received these vitamins during the preconception period experienced a lower incidence of EPL compared to those receiving a placebo. This study suggests that antioxidant supplementation can create a more favorable uterine environment for implantation and early fetal development.⁵²

2. **Coenzyme Q10 (CoQ10):** CoQ10 has been the subject of multiple studies investigating

its role in improving fertility, particularly among women undergoing assisted reproductive technologies (ART). A systematic review of clinical trials indicated that CoQ10 supplementation led to improved ovarian response, increased fertilization rates, and higher pregnancy rates in women undergoing in vitro fertilization (IVF). One study reported that women supplemented with CoQ10 had a significantly lower miscarriage rate compared to those who did not receive the supplement, suggesting that CoQ10 may enhance embryo viability and reduce the risk of EPL.⁵³

3. N-acetylcysteine (NAC): NAC has garnered attention for its antioxidant properties, particularly in women with conditions associated with oxidative stress, such as polycystic ovary syndrome (PCOS). Clinical trials have demonstrated that NAC supplementation can improve insulin sensitivity, ovarian function, and overall reproductive outcomes in women with PCOS. In a specific study, women who received NAC exhibited improved oocyte quality and a reduced rate of EPL compared to a control group. These findings indicate that NAC may provide a protective effect during early pregnancy, particularly in high-risk populations.⁵⁰

4. Combined Antioxidant Therapies: Some clinical evidence suggests that the use of combined antioxidant therapies may offer synergistic benefits. A randomized trial assessed the effects of a multi-antioxidant supplement, including vitamins C and E, CoQ10, and other nutrients, on women with a history of recurrent pregnancy loss. Results indicated that women receiving the combined supplement experienced a statistically significant reduction in EPL rates compared to the control group. This highlights the potential advantage of employing a multi-faceted approach to antioxidant supplementation.⁵¹

5. Dietary Antioxidant Intake: Beyond supplementation, dietary intake of antioxidants has also been linked to reproductive health. A study examining the dietary habits of women trying to conceive found that higher consumption

of antioxidant-rich foods, such as fruits and vegetables, was associated with better reproductive outcomes and a lower risk of miscarriage. This underscores the importance of a balanced diet rich in natural antioxidants in supporting reproductive health and preventing EPL.⁵³

6. Safety and Tolerability: Clinical studies have generally reported that antioxidant supplementation is well-tolerated and safe for most women, even during pregnancy. However, it is crucial to adhere to recommended dosages to avoid potential adverse effects. Monitoring antioxidant levels and assessing the overall health status of patients prior to supplementation can help optimize safety and efficacy.⁵³

Conclusion

Antioxidant supplementation represents a promising strategy for preventing early pregnancy loss (EPL) by addressing the detrimental effects of oxidative stress on reproductive health. The clinical evidence indicates that antioxidants, including vitamins C and E, Coenzyme Q10, and N-acetylcysteine, can improve oocyte and embryo quality, enhance uterine conditions for implantation, and ultimately reduce the risk of EPL. These findings underscore the importance of maintaining a balanced oxidative state during early pregnancy, particularly for women at higher risk of complications due to conditions such as polycystic ovary syndrome or advanced maternal age. The integration of antioxidant supplementation into clinical practice may provide a supportive approach to reproductive health, particularly for women with a history of miscarriage or those undergoing assisted reproductive technologies. However, it is essential for healthcare providers to tailor supplementation regimens to individual needs, considering factors such as medical history and dietary intake.

References

1. Okamgba OC, Nwosu DC, Nwobodo EI, Agu GC, Ozims SJ, Obeagu EI, Ibanga IE, Obioma-Elemba IE, Ihekaire DE, Obasi CC,

- Amah HC. Iron Status of Pregnant and Post-Partum Women with Malaria Parasitaemia in Aba Abia State, Nigeria. *Annals of Clinical and Laboratory Research*. 2017;5(4):206.
- Agreen FC, Obeagu EI. Anaemia among pregnant women: A review of African pregnant teenagers. *Journal of Public Health and Nutrition*. 2023;6(1):138.
 - Obeagu EI, Obeagu GU. Eosinophil Dynamics in Pregnancy among Women Living with HIV: A Comprehensive Review. *Int. J. Curr. Res. Med. Sci.* 2024;10(1):11-24.
 - Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. Evaluation of protein C, protein S and fibrinogen of pregnant women with malaria in Owerri metropolis. *Madonna University journal of Medicine and Health Sciences* ISSN: 2814-3035. 2022 Apr 19;2(2):1-9.
 - Obeagu EI, Obeagu GU. Eosinophilic Changes in Placental Tissues of HIV-Positive Pregnant Women: A Review. *Elite Journal of Laboratory Medicine*, 2024; 2(1): 14-32
 - Joo EH, Kim YR, Kim N, Jung JE, Han SH, Cho HY. Effect of endogenic and exogenic oxidative stress triggers on adverse pregnancy outcomes: preeclampsia, fetal growth restriction, gestational diabetes mellitus and preterm birth. *International journal of molecular sciences*. 2021;22(18):10122.
 - Juan CA, Pérez de la Lastra JM, Plou FJ, Pérez-Lebeña E. The chemistry of reactive oxygen species (ROS) revisited: outlining their role in biological macromolecules (DNA, lipids and proteins) and induced pathologies. *International journal of molecular sciences*. 2021;22(9):4642.
 - Feng Y, Feng Q, Qu H, Song X, Hu J, Xu X, Zhang L, Yin S. Stress adaptation is associated with insulin resistance in women with gestational diabetes mellitus. *Nutrition & diabetes*. 2020;10(1):4.
 - Obeagu EI, Abdirahman BF, Bunu UO, Obeagu GU. Obstetrics characteristics that effect the newborn outcomes. *Int. J. Adv. Res. Biol. Sci.* 2023;10(3):134-43.
 - Anyiam AF, Obeagu EI, Obi E, Omosigho PO, Irondi EA, Arinze-Anyiam OC, Asiyah MK. ABO blood groups and gestational diabetes among pregnant women attending University of Ilorin Teaching Hospital, Kwara State, Nigeria. *International Journal of Research and Reports in Hematology*. 2022;5(2):113-121.
 - Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med.* 2018;3(9):1-4.
 - Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of coagulation parameters in malaria infected pregnant women in Imo state, Nigeria. *International Journal of Current Research in Medical Sciences*. 2018;4(9):41-9.
 - Obeagu EI, Obeagu GU. Neonatal Outcomes in Children Born to Mothers with Severe Malaria, HIV, and Transfusion History: A Review. *Elite Journal of Nursing and Health Science*, 2024; 2(3): 38-58
 - Obeagu EI, Obeagu GU. The Vital Role of Antioxidants in Enhancing Fertility and Pregnancy Success: A Review. *Elite Journal of Nursing and Health Science*. 2023;1(1):1-2.
 - Obeagu EI, Ubosi NI, Uzoma G. Antioxidant Supplementation in Pregnancy: Effects on Maternal and Infant Health. *Int. J. Adv. Multidiscip. Res.* 2023;10(11):60-70.
 - Obeagu EI, Obeagu GU. Enhancing Maternal and Fetal Well-being: The Role of Antioxidants in Pregnancy. *Elite Journal of Medical Sciences*. 2024;2(4):76-87.
 - Obeagu EI, Obeagu GU. Harnessing the Power of Antioxidant-Rich Diet for Preconception Health: A Review. *Elite Journal of Health Science*. 2023;1(1):1-3.
 - Nowak D, Gośliński M, Wojtowicz E, Przygoński K. Antioxidant properties and phenolic compounds of vitamin C-rich juices. *Journal of Food Science*. 2018;83(8):2237-2246.
 - Obeagu EI, Adias TC, Obeagu GU. Influence of Antioxidants on Maternal and Fetal Immune Response: A Review. *Elite Journal of Nursing and Health Science*. 2024;2(6):1-3.
 - Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Neurodevelopmental Outcomes in Offspring: A Review of Maternal

- Interventions. *Elite Journal of Health Science*. 2023;2(5):1-9.
21. Obeagu EI, Batisani K, Obeagu GU. Antioxidants and Postpartum Complications: Preventions. *Elite Journal of Nursing and Health Science*. 2024;2(5):30-40.
 22. Obeagu EI, Obeagu GU. Antioxidants and Gestational Diabetes Mellitus: A Comprehensive Review of Preventive Strategies. *Elite Journal of Health Science*. 2024;2(5):19-29.
 23. Obeagu EI, Obeagu GU. Harnessing the Power of Antioxidants: Enhancing Gamete Quality and Fostering Successful Pregnancy. *Elite Journal of Nursing and Health Science*. 2024;2(3):73-83.
 24. Obeagu EI, Muhimbura E, Obeagu GU. Hypoxia-Induced Oxidative Stress: Maternal and Fetal Implications. *Elite Journal of Haematology*, 2024; 2 (8):57-72.
 25. Obeagu EI, Obeagu GU. Managing Hypoxia in Pregnancy: Current Strategies and Future Directions. *Elite Journal of Medical Sciences*. 2024;2(8):53-63.
 26. Obeagu EI, Obeagu GU. Hypoxia-induced Metabolic Changes in Pregnancy: Clinical Perspectives. *Elite Journal of Medicine*. 2024;2(8):50-9.
 27. Obeagu EI, Chukwu PH. Maternal Well-being in the Face of Hypoxia during Pregnancy: A Review. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2024;11(7):25-38.
 28. Sanchez-Aranguren L, Nadeem S. Bioenergetics adaptations and redox homeostasis in pregnancy and related disorders. *Molecular and Cellular Biochemistry*. 2021;476(11):4003-4018.
 29. Obeagu EI, Obeagu GU. Oxygen Deprivation in Pregnancy: Understanding Hypoxia's Impact on Maternal Health. *Journal home page: <http://www.journalijar.com>*;12(01).
 30. Obeagu EI, Obeagu GU. Hypoxia-Induced Inflammation: Implications for Maternal Health. *Elite Journal of Scientific Research and Review*. 2024;2(6):8-25.
 31. Obeagu EI, Obeagu GU. Hypoxia in Pregnancy: Implications for Fetal Development. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2024;11(7):39-50.
 32. Obeagu EI, Obeagu GU. Hypoxia and Pregnancy: The Role of Genetics and Epigenetics. *Elite Journal of Medical Sciences*. 2024;2(8):24-36.
 33. Carter AM. Evolution of placental function in mammals: the molecular basis of gas and nutrient transfer, hormone secretion, and immune responses. *Physiological reviews*. 2012;92(4):1543-1576.
 34. Obeagu EI, Obeagu GU. Maternal Hypoxia: Impact on Immune System Development in Offspring. *Elite Journal of Health Science*. 2024;2(8):45-57.
 35. Obeagu EI, Obeagu GU. Maternal Hypoxia and Placental Dysfunction: Insights from Molecular Biology. *Elite Journal of Health Science*. 2024;2(8):58-69.
 36. Kalagiri RR, Carder T, Choudhury S, Vora N, Ballard AR, Govande V, Drever N, Beeram MR, Uddin MN. Inflammation in complicated pregnancy and its outcome. *American journal of perinatology*. 2016;33(14):1337-1356.
 37. Al-Gubory KH. Environmental pollutants and lifestyle factors induce oxidative stress and poor prenatal development. *Reproductive biomedicine online*. 2014;29(1):17-31.
 38. Boeldt DS, Bird IM. Vascular adaptation in pregnancy and endothelial dysfunction in preeclampsia. *The Journal of endocrinology*. 2017;232(1):R27.
 39. Burton GJ, Cindrova-Davies T, wa Yung H, Jauniaux E. Hypoxia and reproductive health: Oxygen and development of the human placenta. *Reproduction*. 2021;161(1):F53-65.
 40. He L, He T, Farrar S, Ji L, Liu T, Ma X. Antioxidants maintain cellular redox homeostasis by elimination of reactive oxygen species. *Cellular Physiology and Biochemistry*. 2017;44(2):532-553.
 41. Ighodaro OM, Akinloye OA. First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant defence grid. *Alexandria journal of medicine*. 2018;54(4):287-293.
 42. Roy Z, Bansal R, Siddiqui L, Chaudhary N. Understanding the role of free radicals and antioxidant enzymes in human diseases.

- Current Pharmaceutical Biotechnology. 2023;24(10):1265-1276.
43. Mironczuk-Chodakowska I, Witkowska AM, Zujko ME. Endogenous non-enzymatic antioxidants in the human body. *Advances in medical sciences*. 2018;63(1):68-78.
44. Sebastiani G, Navarro-Tapia E, Almeida-Toledano L, Serra-Delgado M, Paltrinieri AL, García-Algar Ó, Andreu-Fernández V. Effects of antioxidant intake on fetal development and maternal/neonatal health during pregnancy. *Antioxidants*. 2022;11(4):648.
45. Rumbold AR, Crowther CA, Haslam RR, Dekker GA, Robinson JS. Vitamins C and E and the risks of preeclampsia and perinatal complications. *New England Journal of Medicine*. 2006;354(17):1796-1806.
46. Cederberg J, Simán CM, Eriksson UJ. Combined treatment with vitamin E and vitamin C decreases oxidative stress and improves fetal outcome in experimental diabetic pregnancy. *Pediatric research*. 2001;49(6):755-762.
47. Perkins AV. Placental oxidative stress, selenium and preeclampsia. *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health*. 2011;1(1):95-99.
48. Rayman MP, Searle E, Kelly L, Johnsen S, Bodman-Smith K, Bath SC, Mao J, Redman CW. Effect of selenium on markers of risk of pre-eclampsia in UK pregnant women: a randomised, controlled pilot trial. *British Journal of Nutrition*. 2014;112(1):99-111.
49. Luo J, Wu W, Zhang P, Chen X, Feng Y, Ma N, Yang H, Wang Y, Li M, Xie B, Guo P. Zinc levels and birth weight in pregnant women with gestational diabetes mellitus: a matched cohort study in China. *The Journal of Clinical Endocrinology & Metabolism*. 2020;105(7): e2337-2345.
50. Sley EG, Rosen EM, van 't Erve TJ, Sathyanarayana S, Barrett ES, Nguyen RH, Bush NR, Milne GL, Swan SH, Ferguson KK. Omega-3 fatty acid supplement use and oxidative stress levels in pregnancy. *PloS one*. 2020;15(10): e0240244.
51. Orhan H, Önderoglu L, Yücel A, Sahin G. Circulating biomarkers of oxidative stress in complicated pregnancies. *Archives of gynecology and obstetrics*. 2003; 267:189-195.
52. Barbosa ML, de Meneses AA, de Aguiar RP, e Sousa JM, Cavalcante AA, Maluf SW. Oxidative stress, antioxidant defense and depressive disorders: a systematic review of biochemical and molecular markers. *Neurology, Psychiatry and Brain Research*. 2020; 36:65-72.
53. Di Fabrizio C, Giorgione V, Khalil A, Murdoch CE. Antioxidants in pregnancy: do we really need more trials? *Antioxidants*. 2022 Apr 22;11(5):812.

Access this Article in Online



Website:

www.ijcrps.com

Subject:

[Health Sciences](#)

Quick Response Code

DOI: [10.22192/ijcrps.2024.11.09.004](https://doi.org/10.22192/ijcrps.2024.11.09.004)

How to cite this article:

Emmanuel Ifeanyi Obeagu and Getrude Uzoma Obeagu. (2024). Antioxidant Supplementation and Prevention of Early Pregnancy Loss: A Narrative Review. *Int. J. Curr. Res. Chem. Pharm. Sci.* 11(9): 28-37. DOI: <http://dx.doi.org/10.22192/ijcrps.2024.11.09.004>